

DE FREITAS-DOS SANTOS, P. P. (ME)¹, GIMBA, E. R. P.^{1, 2}

¹Grupo de Oncobiologia Celular e Molecular, Coordenação de Pesquisa, Instituto Nacional de Câncer (INCA); ²Departamento de Ciências da Natureza (RCN), Universidade Federal Fluminense.

INTRODUCTION

Prostate cancer antigen 3 (PCA3) is a prostate specific long noncoding RNA (lncRNA) overexpressed in prostate cancer (PCa) tissues in relation non-neoplastic tissues.

Osteopontin (OPN) and TMPRSS2 (T2) genes are among gene products able to modulate PCa cell survival and controlled by androgen receptor (AR) signaling. TMPRSS2 (T2) is responsive to androge, prostate specific, and the T2:ERG gene fusion is the most common gene rearrangement in the CaP, corresponding to 90% of the fusions present in this tumour.




Figure 01. Osteopontin splicing isoforms. OPNa isoform is the complete splice variant, while OPNb does not contain exon 5 and OPNc lacks exon 4.




Figure 02. TMPRSS2:ERG splicing isoforms. T2:ERG full length variant contain N-terminal domain (NTD), Pointed domain (PNT), DNA-binding domain (DBD) and C-terminal domain (CTD). T2:ERG3 isoform lacks 32 amino acids in NTD, and T2:ERG8 a truncated form lacking DBD/CTD, which was replaced by sequences resulting in the addition of 70 unique (U) amino acids.




Figure 03. Androgen receptor pathway. Our group previously demonstrated by Ferreira, et al 2012, that PCA3 expression is regulated by the androgen receptor (AR) signaling. Also, we found that PCA3 knockdown significantly inhibited the expression of AR target genes, besides negatively modulating PCa cell survival.

OBJECTIVES

- Investigate whether PCA3 can modulate the expression of OPN and the T2:ERG fusion splicing isoforms;
- Evaluate how these variants can be associated to PCa cell survival.

METHODOLOGY




Figure 04. The expression of OPN and T2:ERG isoforms have been analyzed by quantitative real time PCR (qRT-PCR). For knockdown assays, the LNCaP prostate cancer cell were cultured in RPMI 1640 culture medium supplemented with 1% FBS without antibiotics. Cells were transfected with a siRNA targeting PCA3 (siPCA3) and a control scrambled siRNA sequence (siSCRBL). Total RNA was extracted using Qiagen Mini kit and cDNA synthesis was done with Superscript II First-Strand Synthesis system. Then, cDNA samples were analyzed by qRT-PCR in order, to validate the PCA3 knockdown. Then, the expression of OPN and T2:ERG isoforms have also been analyzed.

RESULTS




Figure 05. Effect of siPCA3 knockdown on LNCaP cell viability. A) We evaluate the effect of siPCA3 knockdown after 24h and 36h of cell transfection. Transcriptional evaluation of OPN and T2:ERG splicing isoforms have been performed in 24h after siRNA transfection. B) In response to LNCaP transfection with siPCA3 there was a significant decrease on cell viability, when compared to siSCRBL-transfected cells.




Figure 06. Relative expression levels of OPN isoforms in response to PCA3 knockdown. The transcriptional levels of the OPN isoforms were analyzed by qRT-PCR, using oligonucleotides specific for these isoforms and GAPDH was used as constitutive expression gene. The results presented were performed in duplicate and are representative of 1 assay, using the siSCRBL as the reference sample (reference value = 1). In response to PCA3 knockdown, we observed a significant decreased on the expression of the three OPN isoforms.




Figure 07. Relative expression levels of T2:ERG isoforms in response to PCA3 knockdown. The transcriptional levels of the T2:ERG isoforms were analyzed by qRT-PCR, using oligonucleotides specific for these isoforms and GAPDH was used as constitutive expression gene. The results presented were performed in duplicate and are representative of 3 independent assays, using the T2:ERG as reference sample (reference value = 1). In response to PCA3 knockdown, we observed a significant decrease on the expression levels of the three T2:ERG isoforms. The P values refer to the comparison between the level of transcriptional expression of T2:ERG and T2:ERG3 and T2:ERG8 isoforms. *** P < 0.001.

CONCLUSIONS

We found that in response to PCA3 knockdown, all three OPN isoforms significantly decreased their expression levels, as well as all the three T2:ERG variants. Among these isoforms, OPNc and the T2:ERG3 variants were most downregulated. These data provide early evidence regarding the putative role of PCA3 as modulator of T2:ERG and OPN splicing isoforms and that PCA3 roles on modulating survival can involve the gene target.

FINANCIAL SUPPORT: •CNPq, CAPES, FAPERJ, INCT do Câncer, Ministério da Saúde, Proppi-UFF.

Projeto Gráfico: Setor de Edição e Informação Técnico-Científica / INCA